

The Early Life History of the Clam *Macoma balthica* in a High CO₂ World

Carl Van Colen*, Elisabeth Debusschere, Ulrike Braeckman, Dirk Van Gansbeke, Magda Vincx

Ghent University, Department of Biology, Marine Biology Section, Ghent, Belgium

Abstract

This study investigated the effects of experimentally manipulated seawater carbonate chemistry on several early life history processes of the Baltic tellin (*Macoma balthica*), a widely distributed bivalve that plays a critical role in the functioning of many coastal habitats. We demonstrate that ocean acidification significantly depresses fertilization, embryogenesis, larval development and survival during the pelagic phase. Fertilization and the formation of a D-shaped shell during embryogenesis were severely diminished: successful fertilization was reduced by 11% at a 0.6 pH unit decrease from present (pH 8.1) conditions, while hatching success was depressed by 34 and 87%, respectively at a 0.3 and 0.6 pH unit decrease. Under acidified conditions, larvae were still able to develop a shell during the post-embryonic phase, but higher larval mortality rates indicate that fewer larvae may metamorphose and settle in an acidified ocean. The cumulative impact of decreasing seawater pH on fertilization, embryogenesis and survival to the benthic stage is estimated to reduce the number of competent settlers by 38% for a 0.3 pH unit decrease, and by 89% for a 0.6 pH unit decrease from present conditions. Additionally, slower growth rates and a delayed metamorphosis at a smaller size were indicative for larvae developed under acidified conditions. This may further decline the recruit population size due to a longer subjection to perturbations, such as predation, during the pelagic phase. In general, early life history processes were most severely compromised at ~pH 7.5, which corresponds to seawater undersaturated with respect to aragonite. Since recent models predict a comparable decrease in pH in coastal waters in the near future, this study indicates that future populations of *Macoma balthica* are likely to decline as a consequence of ongoing ocean acidification.

Citation: Van Colen C, Debusschere E, Braeckman U, Van Gansbeke D, Vincx M (2012) The Early Life History of the Clam *Macoma balthica* in a High CO₂ World. PLoS ONE 7(9): e44655. doi:10.1371/journal.pone.0044655

Editor: Patrick Callaerts, VIB & Katholieke Universiteit Leuven, Belgium

Received: November 24, 2011; **Accepted:** August 10, 2012; **Published:** September 10, 2012

Copyright: © 2012 Van Colen et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: CVC is a postdoctoral fellow of the Flemish Fund for Scientific Research FWO (1.2.380.11.N). Additional financial support for this study came from the special research fund of Ghent University through project BOF-GOA 01GA1911W. The funder had no role in study design, data collection and analysis.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: carl.vancolen@ugent.be

Introduction

About one third of the total anthropogenic CO₂ emissions has currently been absorbed by the oceans, driven by the difference of pCO₂ in the atmosphere and the surface ocean layers [1]. This results in a continuous increase in seawater pCO₂ (i.e. hypercapnia) and a decrease in seawater pH and carbonate ion (CO₃²⁻) availability [2], in a process referred to as ocean acidification. The pH of seawater has already declined by 0.1 unit compared to pre-industrial values and several climate change models predict further decreases up to 0.35 units by the end of this century [3,4]. However, recent studies indicate that the pH in temperate coastal systems is likely to decrease and order of magnitude faster due to an altered balance between primary production and respiration [5,6]. These fast-occurring changes may pose far-reaching consequences for marine ecosystems since empirical evidence demonstrates significant alterations in trophodynamics, nutrient cycling, organism physiology, organism reproduction and development as a consequence of ocean acidification (reviewed in e.g. [7]). The implications of ocean acidification for ecosystem resilience are, however, still debated (see e.g. [8–10]) because the observed responses are variable and it remains unclear how acidification will interact with other stressors, such as temperature rise, eutrophication and deoxygenation of the oceans.

Meta-analyses and literature reviews suggest that particularly calcification processes are hampered by ocean acidification, e.g. [4,8,10,11]. Subsequently, calcifying organisms are considered specifically susceptible to ocean acidification. The formation of calcium carbonate structures mainly depend on the availability of carbonate ions, given the fact that (1) Ca²⁺ concentrations are rather constant at a given salinity, and (2) the CaCO₃ saturation state (Ω) is defined as:

$$\Omega = \frac{[CO_3^{2-}][Ca^{2+}]}{K'_{sp}};$$

where K'_{sp} is the stoichiometric solubility product which depends on temperature, salinity, pressure and the mineral phase considered (e.g. high-magnesian calcite, calcite, aragonite). In an acidified ocean, fewer CO₃²⁻ ions become available to form CaCO₃ structures and reduced calcification rates have been observed in a variety of calcifying organisms (see e.g. [12] for a review). However, calcification responses to ocean acidification may vary amongst organisms and between life stages, depending on species-specific life-history traits. For example, some organisms may be able to compensate for alteration in carbonate chemistry by increasing calcification rates [13], and organisms which have sufficient energy reserves may be more resilient to ocean

acidification [14]. In general, early life history stages are believed to be particularly vulnerable to decreases in seawater pH because CaCO₃ mineral phases of early life stages (e.g. amorphous CaCO₃ and aragonite) are often less stable and more soluble than the CaCO₃ mineral phases which are predominantly precipitated by adults (e.g. calcite) [15–17]. Consequently, most ocean acidification studies on the early development of a variety of invertebrate phyla reveal inhibitive effects on larval fitness along with decreasing seawater pH: decreased growth, increased mortality and malformations, e.g. [18–24].

Understanding ocean acidification effects on early life history processes is crucial in assessing future ecosystem structure and functioning because settlement and recruitment are fundamental processes in driving ecological balance of communities. This study investigated whether ocean acidification affects several early developmental processes of the Baltic tellin (*Macoma balthica*), from fertilization until metamorphosis. *Macoma balthica* is known to influence sediment nutrient cycling [25,26] and sediment erodibility in coastal habitats [27]. Furthermore *M. balthica* is a key species in ecosystem carbon flow, because it is a dominant grazer of microalgae and a favourable food source for wading birds, epibenthic crustaceans and fishes [28–30]. This species occurs from the Gironde estuary in Southwest France to the polar region in Greenland and Siberia and on the east and west coasts of North America (www.iobis.org), where it often constitutes significant amounts of the benthic metazoan biomass, e.g. [31,32]. Given the wide distribution of the species and its importance to ecosystem processes, the present study provides a significant contribution to the assessment of future ecosystem functioning in the light of ongoing acidification in coastal and estuarine waters.

Materials and Methods

Production of Eggs, Embryos and Larvae: General Procedures

About 600 adult individuals were collected during low tide in April 2011 at the Paulinaschor, a tidal flat located in the polyhaline part of the Westerschelde estuary (51°21'24"N, 3°42'51"E; permits for the field work were approved and obtained by Provincie Zeeland, the Netherlands; Directie Ruimte, Milieu en Water). Test organisms were stored in the lab in containers (40×33×14 cm; about 150 individuals per container) filled with sieved sediment (1 mm) and aerated 25 PSU filtered seawater at 5°C. They were fed three times a week on a mixed diet of *Isochrysis*, *Pavlova*, *Thalassiosira weissflogii* and *Tetraselmis* (Shellfish Diet 1800; Reed Mariculture). Individual organisms were induced to spawn in 25 PSU 0.2 µm filtered seawater by exposure to the selective serotonin re-uptake inhibitor fluoxetine, preceded by a Δ 10°C temperature shock [33]. On average, 22.3% of the adults could be induced to spawn. Fertilization was carried out by pipetting eggs of several females into a clean glass beaker and adding 1 to 3 ml of a homogenized sperm suspension derived from several males. The resultant mixture was left undisturbed for 4 h at 15°C. Embryos (diameter ~100 µm) were then separated from all other matter by rinsing them over stacked sieves of 125 and 32 µm. Subsequently, they were transferred into 2.3 L culture flasks containing 15°C UV-irradiated filtered seawater with a salinity of 25 and dosed with 1.5×10⁻⁵ g.L⁻¹ Penicillin G potassium salt and 2.5×10⁻⁵ g.L⁻¹ streptomycin sulphate (further referred to as FSW-AB). The culture flasks were placed on a roller-table (3 rpm) to avoid sinking of embryos and larvae.

Carbonate Chemistry Manipulation and Measurements

Filtered seawater was diluted with tap water until 25 PSU, i.e. the average salinity at the site where adult organisms were retrieved, and incubated at 15°C and UV-irradiated for 1 week prior to the use in all experiments. Seawater carbonate chemistry was manipulated through direct bubbling of seawater (~pH 8.1) with 100% CO₂ gas using Dulcometer technology (ProMinent®) in order to create predicted pH conditions for likely outcomes within the current century in coastal waters: ~pH 7.8, and ~pH 7.5 [6,34]. This method was chosen as it best replicates ocean acidification by altering dissolved inorganic carbon while keeping total alkalinity constant [35]. The environmental parameters and carbonate chemistry conditions that persisted during each experiment are outlined in Tables S1, S2, and S3.

The carbonate chemistry manipulation unit comprised two incubation tanks (60 L and 15 L) for each pH treatment. Seawater was pumped constantly between the two tanks in order to mix the seawater (5 L.min⁻¹) and one third of the total volume was replaced every day by freshly incubated seawater. The 15 L incubation tank was constantly bubbled with ambient air to maintain dissolved oxygen levels >90%. The carbonate chemistry was manipulated through controlled pumping of 100% CO₂ gas at the bottom of the 60 L incubation tank, using a pH controller (Dulcometer D1C, version B) connected to a pH electrode (Dulcotest PHE-112SE) mounted in the 60 L incubation tank. Every other day the pH electrodes were calibrated using Hanna Instruments' pH_{NBS} buffers. The control seawater unit had the same set-up except for the CO₂ gas supply and was bubbled with ambient air. Seawater with the targeted pH was pumped from the manipulation units directly into the culture flasks without leaving headspace to avoid water-atmosphere gas exchange [35]. Temperature and salinity (WTW COND 330), pH (Dulcotest PHE-112SE) and total alkalinity (TA) were measured prior and at the end of each incubation in the different experiments, as further described below. Total alkalinity was determined on 20 ml samples filtered on GF/F membranes by end-point titration of 25 mmol.L⁻¹ HCl, using a Mettler Toledo G20 Compact Titrator. The recorded values were entered in CO2SYS software [36] to calculate the remaining carbonate chemistry parameters using the thermodynamic constants of Merzbach et al. [37]: partial CO₂ pressure, total concentration of dissolved inorganic carbon (DIC), concentration of HCO₃⁻ and CO₃²⁻, saturation state of aragonite and calcite. We are aware that the applied method for alkalinity determination differs from the general procedure applied in the ocean acidification community (Gran Titration, [38]); however our measurements of Dickson CRM water (batch 107) had, on average, only 3.35% higher TA values. Since these enhanced TA values resulted in only slightly overestimated concentrations of CO₃²⁻ (i.e. on average +20.7, +10.6, +6.4 µmol.kg⁻¹ for control, ~pH_{7.8} and ~pH_{7.5} conditions) and aragonite saturation states (i.e. on average +0.33, +0.17, +0.10 for control, ~pH_{7.8} and ~pH_{7.5} conditions) we are confident that the obtained results and conclusions deduced from this study are not compromised.

Ocean Acidification Effects on Fertilization Success

Fertilization was carried out by pipetting 50 eggs into clean vials containing 23 ml of control, ~pH_{7.8} and ~pH_{7.5} seawater. Subsequently, 2 ml of the sperm suspension was added. Five replicate vials of each pH treatment were continuously and gently shaken during incubation at 15.2°C for 24 hours. Embryonic development was stopped by adding 1 ml of a neutralized 4% formaldehyde – tap water solution. Unfertilized eggs and the embryos already developed were counted using a Leica MZ 16

Table 1. Analysis of variance on difference in early life history processes of *Macoma balthica* reared in the different pH treatments, and correlation with seawater pH.

Developmental parameter	Test statistic	p-level	Post-hoc tests	Correlation with pH
Fertilization success	F = 9.581	0.003	control ≠ ~pH _{7.5} ; ~pH _{7.8} ≠ ~pH _{7.5}	r = 0.71, p = 0.003**
Hatching success	H = 7.200	0.027	control ≠ ~pH _{7.8} ≠ ~pH _{7.5}	r = 0.97, p < 0.001*
Shell size at day 3	H = 7.200	0.027	control ≠ ~pH _{7.8} ≠ ~pH _{7.5}	r = 0.94, p < 0.001*
Growth rate	H = 7.200	0.027	control ≠ ~pH _{7.8} ≠ ~pH _{7.5}	r = 0.93, p < 0.001**
Mortality rate	H = 6.489	0.039	control ≠ ~pH _{7.5}	r = -0.82, p = 0.007*
% of larvae metamorphosed at day 19	H = 1.689	0.430		r = -0.01, p = 0.840*

*indicate results from product moment correlations of logarithmic transformed data,

**indicate results from non-parametric Spearman-rank correlation. For more details: see text.

doi:10.1371/journal.pone.0044655.t001

stereomicroscope and the fertilization success was calculated as:

$$\text{Fertilization success} = \frac{\text{developed embryos}}{(\text{unfertilized eggs} + \text{developed embryos})}$$

Ocean Acidification Effects on Embryonic Development

Adults were induced to spawn and eggs were fertilized under control conditions according to the general procedures described above. Subsequently, embryos were equally distributed into triplicated 2.3 L culture flasks (2.2 embryos ml⁻¹) containing 15.2°C control, ~pH_{7.8} and ~pH_{7.5} FSW-AB and left to incubate on a roller-table. After 3 days, five 4 ml aliquots were collected from each culture flask and embryonic development was stopped by adding 1 ml of a neutralized 4% formaldehyde – tap water solution to the aliquot. The shell length of all embryos that had developed a shell were measured to the nearest 0.1 μm under a stereomicroscope using a camera lucida mounted to a stereomicroscope and Image J 1.44 software. In addition, the density of unfertilized eggs, embryos that have developed to a normal D-shaped shell characterized by a straight hinge, and abnormal developed larvae (i.e. shell with a convex hinge or protruding mantle; [39]) was determined in order to calculate hatching success according to:

$$\text{Hatching success} = \frac{\text{D-shaped larvae}}{(\text{unfertilized eggs} + \text{abnormal developed larvae} + \text{D-shaped larvae})}$$

Ocean Acidification Effects on Larval Development

Adults were induced to spawn and embryos were developed under control conditions according to the general procedures described above. Subsequently, 3 day-old larvae were equally distributed into triplicated 2.3 L culture flasks (1.4 larvae.ml⁻¹) containing 14.8°C control, ~pH_{7.8} and ~pH_{7.5} FSW-AB and 10⁵ live *Isochrysis galbana* cells.ml⁻¹ originating from a continuously maintained healthy culture in the laboratory. The larval cultures were incubated on a roller-table for 16 days (i.e. in total 19 days after fertilization) and refreshed every other day by rinsing the seawater and larvae over a 32 μm mesh-sized sieve and transferring the larvae into new culture flasks containing control,

~pH_{7.8} and ~pH_{7.5} FSW-AB and 10⁵ live *I. galbana* cells.ml⁻¹. Seawater subsamples (25 ml) were collected from each culture flask at the start and end of incubation in order to calculate the carbonate chemistry parameters. Additionally, three subsamples were taken every two days from each culture flask to measure larval mortality, growth and metamorphosis stage. Following Bos et al. [40,41] we obtained:

- (1) mortality rates from fitting a regression line through the natural logarithms of the densities through time, according to:

$$\ln(N_t) = \ln(N_0) + rt;$$

where N_t is the density of larvae at time t (larvae.ml⁻¹), N_0 the initial density at day 3 (larvae.ml⁻¹), r is the mortality rate (d⁻¹), t is the age (d) of the larvae;

- (2) growth rates from the linear model:

$$SL = SL_3 + b(\text{Age} - 3);$$

where SL is the shell length (μm), SL₃ is the initial shell length (μm) at day 3 and b is the growth rate (μm.d⁻¹);

- (3) average shell length at metamorphosis from the logistic regression:

$$\ln[p(1-p)^{-1}] = \alpha + bSL;$$

where p is the probability of passing metamorphosis (i.e. the presence of a clearly visible foot bulging out of the shell). The age at which 50% of the larvae had metamorphosed (i.e. reaction norm) was calculated by setting p at 0.5 in the logistic regression model; yielding the linear predictor, $lp=0$. The average age at which 50% of the larvae reach metamorphosis was subsequently calculated from the modified equation (2):

$$\text{Age} = 3 + \frac{(SL - SL_3)}{b};$$




Figure 1. *Macoma balthica*: Ocean acidification effects on fertilization and embryogenesis. (a) fertilization success, (b) hatching success, and (c) size of larvae that have developed a shell after 3 days, reared at three pH conditions: control (\sim pH_{8.1}), \sim pH_{7.8} and \sim pH_{7.5}. Level of replication: $n=5$ for fertilization success, $n=3$ for hatching success and size. Presented error bars are standard errors. doi:10.1371/journal.pone.0044655.g001

where SL is the shell length at metamorphosis deduced from the logistic regression.

Data Analysis

One-way analysis of variance, followed by Tukey HSD post-hoc tests, was used to interpret the significance of ocean acidification effects on fertilization success ($n=5$), after testing for normality (Shapiro-Wilk's test) and homoscedasticity (Levene test). Kruskal-Wallis rank sum tests followed by Conover's multiple comparison tests [42] were applied to investigate ocean acidification effects on hatching success, the percentage of metamorphosed larvae at day 19, larval mortality and growth rates ($n=3$). For all tests,

differences were considered significant at $p<0.05$. In order to interpret relationships of developmental parameters with the manipulated seawater carbonate chemistry we performed correlations with pH. In addition, Michaelis-Menten functions were fitted to the CO_3^{2-} ion concentration since the latter parameter is considered the main governing factor of larval shellfish growth [23].

Results

Fertilization and Embryonic Development

Manipulation of seawater carbonate chemistry significantly altered egg fertilization and embryonic development to a D-shaped larvae (Table 1). On average, only 76.19 ± 0.02 SE % of the eggs were successfully fertilized at \sim pH_{7.5}, which was respectively 9 and 8% lower as compared to fertilization under control and \sim pH_{7.8} conditions (Fig. 1a). In addition, significantly fewer embryos were able to produce a normal D-shape after 3 days of development at \sim pH_{7.8} and \sim pH_{7.5} as compared to control embryos (Fig. 1b). Hatching success for embryos reared at \sim pH_{7.5} were 48 and 29% lower as compared to embryos reared under control and \sim pH_{7.8} conditions, respectively. On average, 3 day-old control embryos that have developed a shell measured 155.4 ± 0.4 SE μm , which was significantly larger than embryos reared under acidified conditions: i.e. shell size was 148.4 ± 1.0 SE μm and 138.7 ± 2.5 SE μm , respectively at \sim pH_{7.8} and \sim pH_{7.5} (Fig. 1c). Simple correlations indicate that fertilization and embryonic development significantly decreased with decreasing pH (Table 1), while non-linear regressions indicate that decreasing CO_3^{2-} concentrations resulted in smaller larvae and a drastically lower hatching success, especially below the aragonite saturation level ($\Omega_{\text{aragonite}}$) (Fig. 2b, c).

Larval Development

Larval development was significantly affected by the carbonate chemistry of the seawater (Table 1). The initial shell length was 156.0 ± 1.0 SE μm at day 3 for all treatments. At day 19, larvae grown under control conditions had an average shell length of 308.4 ± 3.4 SE μm , while larvae grown under acidified conditions had an average shell length of 257.6 ± 3.2 SE μm (\sim pH_{7.8}) and 256.1 ± 2.3 SE μm (\sim pH_{7.5}). The initial density of larvae was 1.36 larvae. ml^{-1} at day 3. Under control conditions, the larval density was reduced by 93% to 0.10 ± 0.01 SE larvae. ml^{-1} at day 19. Mortality of larvae that developed under acidified conditions was even more pronounced (97 ± 0.007 SE % and 99 ± 0.004 SE %, respectively at \sim pH_{7.8} and \sim pH_{7.5}). Thus, larvae grow more slowly and suffer greater mortality under acidified conditions (Fig. 3a, b), as was confirmed by the decrease in growth rate and increase in mortality rates when fewer CO_3^{2-} ions became available (Table 1) (Fig. 2d, e). However, multiple comparison tests only detected significantly different mortality rates between larvae that developed under control conditions and at \sim pH_{7.5}.

There was no significant difference among treatments in the percentage of larvae that had metamorphosed at day 19 (Fig. 3c). However logistic regression indicated that, on average, larvae reared under acidified conditions metamorphosed at a smaller size (267.05 μm and 261.35 μm at \sim pH_{7.8} and \sim pH_{7.5}, respectively) which was reached later in the development process, i.e. after 16.3–16.4 days, as compared to larvae that have developed under control conditions (size at metamorphosis = 281.5 μm , age at metamorphosis = 15.1 days).




Figure 2. *Macoma balthica*: Relationships between seawater carbonate ion concentration [CO₃²⁻] and parameters of early life history processes. Fertilization (upper panel; a), embryogenesis (middle panel; b,c), and larval development (lower panel; d,e). The dashed lines refer to the carbonate concentration at the aragonite saturation level (Ω_{aragonite} = 1). The fitted lines show the relationship between the CO₃²⁻ concentration and the developmental parameter according to a Michaelis-Menten equation, e.g. Hatching success = $V_{max}[\text{CO}_3^{2-}]/(K_m + [\text{CO}_3^{2-}])$; where V_{max} is the maximal hatching success and K_m is the CO₃²⁻ concentration where the reaction rate is the half of V_{max} . Percentage of variation explained (r^2) by the fitted lines are 0.45 (fertilization success), 0.90 (size at day 3), 0.86 (hatching success), 0.64 (larval mortality rate), and 0.87 (larval growth rate). doi:10.1371/journal.pone.0044655.g002

Discussion

This study illustrates that ocean acidification may negatively affect shellfish recruitment success by impacting multiple early life history processes prior to settlement, including egg fertilization, embryonic shell formation (i.e. hatching success), larval mortality, growth and metamorphosis.

Results of the fertilization experiment demonstrate that failure of fertilizations increase with enhanced seawater pCO₂. Similar effects have been found for a variety of invertebrate phyla and have been attributed to a reduction in the efficiency to block polyspermy and a reduction in sperm speed and motility which decreases fertilization success (e.g. [43–46]). In addition, the intracellular egg pH has been shown instrumental to successful fertilization of sea urchin eggs by regulating sperm entrance through the egg membrane [47]. This mechanism may therefore be distorted when more CO₂ diffuses across the gamete cell

membrane in an acidified sea. So far studies of ocean acidification effects on bivalve fertilization have yielded variable results. This suggests that effects may be species-specific and reflect the adaptation of the species to the pH variability in the species' habitat, and that effects may be influenced by site-specific environmental parameters, such as temperature and eutrophication. For example, Parker et al. [48] demonstrated that fertilization in a population of the Pacific oyster *Crassostrea gigas* from Port Stevens (Australia) was significantly reduced at elevated CO₂, while Havenhand & Schlegel [49] did not detect a significant effect for a *C. gigas* population from Western Sweden. We found a significant reduction in fertilization success of *Macoma balthica* at ~pH_{7.5} but not at ~pH_{7.8} which may indicate adaptation to fluctuating pH conditions that characterize the habitat of the studied population, i.e. pH 8.2–7.8 [50].

Under acidified conditions, *Macoma balthica* embryos were still able to form a shell, even in seawater undersaturated with respect




Figure 3. *Macoma balthica*: Ocean acidification effects on larval development. Shell length, survival and development stage of larvae between day 3 and day 19 of development reared at three seawater pH conditions: \sim pH_{8.1} (circles, solid line), \sim pH_{7.8} (squares, dashed line), and \sim pH_{7.5} (triangles, dotted line). Regression lines in (a) and (b) indicate average larval growth rate ($\mu\text{m}\cdot\text{d}^{-1}$) and mortality rate (d^{-1}) for each treatment.

doi:10.1371/journal.pone.0044655.g003

to aragonite. However, larval growth rates were significantly reduced under acidified conditions, which corroborate other research on bivalve early life history development, e.g. [12,18,23,24,51]. Non-linear relationships between seawater CO₃²⁻ concentrations and growth rate and hatching success, with a sharp decrease below the aragonite saturation level, demonstrate the importance of the amount of available CO₃²⁻

ions to form CaCO₃ shells. The observed higher larval mortality and a longer development time of larvae in acidified seawater likely reflects a higher energy demand to form a shell when less CO₃²⁻ ions are available.

Recruitment, which demands successful larval settlement and post-settlement growth and survival is critical to the persistence and resilience of benthic populations. Pre-recruitment mortality of planktonic invertebrate larvae is variable but generally accepted to be very high (3–23% daily), particularly due to predation and environmental factors [52,53]. Ocean acidification thus exerts a significant additional burden to the recruitment of *Macoma balthica* by negatively impacting multiple early life history processes. Assuming that abnormally developed 3 day-old larvae will not develop to the benthic stage, the quantified effects of decreasing seawater pH on fertilization, embryogenesis and larval development in this study suggest that the number of *M. balthica* larvae that are competent to settle will decline up to 38–89%, depending on the scenario (i.e. respectively for a 0.3 and 0.6 pH unit decrease from present conditions). Additionally, a delayed metamorphosis at a smaller larval size may further diminish the recruit population size due to a longer subjection to perturbations (e.g. predation) during the pelagic phase. As *Macoma balthica* is recognized as an ecosystem engineer to coastal habitats, e.g. [26,54], ocean acidification-mediated changes to this species' population size and distribution may affect coastal diversity and ecosystem functioning.

Supporting Information

Table S1 Environmental parameters and carbonate chemistry for the different treatments to investigate ocean acidification effects on fertilization success. Carbonate parameters were computed from the recorded seawater pH_{NBS} and Total Alkalinity during the incubations. - : no data available.

(XLSX)

Table S2 Environmental parameters and carbonate chemistry for the different treatments to investigate ocean acidification effects on embryonic development. Carbonate parameters were computed from the recorded seawater pH_{NBS} and Total Alkalinity during the incubations.

(XLSX)

Table S3 Environmental parameters and carbonate chemistry for the different treatments to investigate ocean acidification effects on larval development. Carbonate parameters were computed from the recorded seawater pH_{NBS} and Total Alkalinity during the incubations. - : no data available.

(XLSX)

Acknowledgments

We are grateful to Emily Dolan and Jeroen Ingels for editing and proofreading the manuscript. We also thank two anonymous reviewers for providing useful comments which improved this manuscript. Stijn Bruwier is acknowledged for discussions on the manipulation of seawater carbonate chemistry through controlled bubbling with CO₂ gas.

Author Contributions

Conceived and designed the experiments: CVC ED DVG. Performed the experiments: CVC ED. Analyzed the data: CVC ED DVG UB. Wrote the paper: CVC ED UB MV.

References

- Sabine CL, Feely RA, Gruber N, Key RM, Lee K, et al. (2004) The oceanic sink for anthropogenic CO₂. *Science* 305: 367–371.
- Feely RA, Doney SC, Cooley SR (2009) Ocean Acidification: Present Conditions and Future Changes in a High CO₂ World. *Oceanography* 22: 36–47.
- Caldeira K, Wickett ME (2003) Anthropogenic carbon and ocean pH. *Nature* 425: 365–365.
- Orr JC, Fabry VJ, Aumont O, Bopp L, Doney SC, et al. (2005) Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* 437: 681–686.
- Wootton JT, Pfister CA, Forester JD (2008) Dynamic patterns and ecological impacts of declining ocean pH in a high-resolution multi-year dataset. *Proceedings of the National Academy of Sciences of the United States of America* 105: 18848–18853.
- Provoost P, van Heuven S, Soetaert K, Laane RWPM, Middelburg JJ (2010) Seasonal and long-term changes in pH in the Dutch coastal zone. *Biogeosciences* 7: 3869–3878.
- Fabry VJ, Seibel BA, Feely RA, Orr JC (2008) Impacts of ocean acidification on marine fauna and ecosystem processes. *Ices Journal of Marine Science* 65: 414–432.
- Hendriks IE, Duarte CM, Alvarez M (2010) Vulnerability of marine biodiversity to ocean acidification: a meta-analysis. *Estuarine, Coastal and Shelf Science* 86: 157–164.
- Dupont S, Dorey N, Thorndyke M (2010) What meta-analysis can tell us about vulnerability of marine biodiversity to ocean acidification. *Estuarine, Coastal and Shelf Science* 89: 182–185.
- Hendriks IE, Duarte CM (2010) Ocean acidification: separating evidence from judgment – a reply to Dupont, et al. *Estuarine, Coastal and Shelf Science* 89: 186–190.
- Kroeker KJ, Kordas RL, Crim RN, Singh GG (2010) Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms. *Ecology letters* 13: 1419–1434.
- Kurihara H (2008) Effects of CO₂-driven ocean acidification on the early developmental stages of invertebrates. *Marine Ecology Progress Series* 373: 275–284.
- Guttowska MA, Pörtner HO, Melzner F (2008) Growth and calcification in the cephalopod *Sepia officinalis* under elevated seawater pCO₂. *Marine Ecology Progress Series* 373: 303–309.
- Dupont S, Lundv B, Thorndyke M (2010) Near future ocean acidification increases growth rate of the lecithotrophic larvae and juveniles of the sea star *Crossaster papposus*. *Journal of Experimental Zoology* 314B: 382–389.
- Mucci A (1983) The Solubility of Calcite and Aragonite in Seawater at Various Salinities, Temperatures, and One Atmosphere Total Pressure. *American Journal of Science* 283: 780–799.
- Weiss IM, Tuross N, Addadi L, Weiner S (2002) Mollusc larval shell formation: amorphous calcium carbonate is a precursor phase for aragonite. *Journal of Experimental Zoology* 293: 478–491.
- Addadi L, Raz S, Wiener S (2003) Taking advantage of disorder: amorphous calcium carbonate and its roles in biomineralization. *Advanced Materials* 15: 959–970.
- Talmage SC, Gobler CJ (2010) Effects of past, present, and future ocean carbon dioxide concentrations on the growth and survival of larval shellfish. *Proceedings of the National Academy of Sciences* 107: 17246–17251.
- Lischka S, Bubenbender J, Boxhammer T, Riebesell U (2010) Impact of ocean acidification and elevated temperatures on early juveniles of the polar shelled pteropod *Limacina helicina*: mortality, shell degradation, and shell growth. *Biogeosciences discussions* 7: 8177–8214.
- Comeau S, Gorsky G, Alliouane S, Gattuso JP (2010) Larvae of the pteropod *Cavolinia inflexa* exposed to aragonite undersaturation are viable but shell-less. *Marine Biology* 157: 2341–2345.
- Ellis RP, Bersey J, Rundle SD, Hall-Spencer JM, Spicer JI (2009) Subtle but significant effects of CO₂ acidified seawater on embryos of the intertidal snail, *Littorina obtusata*. *Aquatic Biology* 5: 41–48.
- Watson SA, Southgate PC, Tyler PA, Peck LS (2009) Early larval development of the Sydney rock oyster *Saccostrea glomerata* under near-future predictions of CO₂-driven ocean acidification. *Journal of Shellfish Research* 28: 431–437.
- Gazeau F, Gattuso JP, Dawber C, Pronker AE, Peene F, et al. (2010) Effect of ocean acidification on the early life stages of the blue mussel *Mytilus edulis*. *Biogeosciences* 7: 2051–2060.
- Gazeau F, Gattuso JP, Greaves M, Elderfield H, Peene J, et al. (2011) Effect of carbonate chemistry alteration on the early embryonic development of the Pacific oyster (*Crassostrea gigas*). *Plos One*, 6, e23010/1–8.
- Marinelli RL, Williams TJ (2003) Evidence for density-dependent effects of infauna on sediment biogeochemistry and benthic-pelagic coupling in nearshore systems. *Estuarine Coastal and Shelf Science* 57: 179–192.
- Michaud E, Desrosiers G, Mermillod-Blondin F, Sundby B, Stora G (2006) The functional group approach to bioturbation: II. The effects of the *Macoma balthica* community on fluxes of nutrients and dissolved organic carbon across the sediment-water interface. *Journal of Experimental Marine Biology and Ecology* 337: 178–189.
- Willows RI, Widdows J, Wood RG (1998) Influence of an infaunal bivalve on the erosion of an intertidal cohesive sediment: A flume and modeling study. *Limnology and Oceanography* 43: 1332–1343.
- Zwarts L, Blomert AM (1992) Why Knot *Calidris canutus* Take Medium-Sized *Macoma balthica* When 6 Prey Species Are Available. *Marine Ecology Progress Series* 83: 113–128.
- Herman PMJ, Middelburg JJ, Widdows J, Lucas CH, Heip CHR (2000) Stable isotopes as trophic tracers: combining field sampling and manipulative labelling of food resources for macrobenthos. *Marine Ecology-Progress Series* 204: 79–92.
- Hiddink JG, Marijnissen SAE, Troost K, Wolff WJ (2002) Predation on O-group and older year classes of the bivalve *Macoma balthica*: interaction of size selection and intertidal distribution of epibenthic predators. *Journal of Experimental Marine Biology and Ecology* 269: 223–248.
- Holland AF, Shaughnessy AT, Hiegel MH (1987). Long-term variation in mesohaline Chesapeake Bay macrobenthos: spatial and temporal patterns. *Estuaries* 10: 227–245.
- Van Colen C, Monserrat F, Vincx M, Herman PMJ, Ysebaert T, et al. (2010) Long-term divergent tidal flat benthic community recovery following hypoxia-induced mortality. *Marine Pollution Bulletin* 60: 178–186.
- Honkoop PJC, Luttikhuisen PC, Piersma T (1999) Experimentally extending the spawning season of a marine bivalve using temperature change and fluoxetine as synergistic triggers. *Marine Ecology Progress Series* 180: 297–300.
- IPCC: Climate Change 2007: The physical science basis (2007). Contribution of working group A to the fourth assessment. Report of the Intergovernmental Panel on Climate Change (eds Solomon S, Qin D, Manning M, Chen Z, Marquis M, et al.). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 996 p.
- Gattuso JP, Gao K, Lee K, Rost B, Schulz KG (2010) Approaches and tools to manipulate the carbonate chemistry. In: *Guide for best practices in ocean acidification research and data reporting* (eds Riebesell U, Fabry V, Hansson L, Gattuso JP), Office for Official Publications of the European Communities, Luxembourg.
- Pelletier G, Lewis E, Wallace D (2007) CO₂SYSL: a calculator for the CO₂ system in seawater for Microsoft Excel/VBA, Washington State Department of Ecology/Brookhaven National Laboratory, Olympia, WA/Upton, NY, USA.
- Merzbach C, Culbertson CH, Hawley JE, Pytkowicz RM (1973) Measurements of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnology and Oceanography* 18: 897–907.
- Dickson AG, Sabine CL, Christian JR (2007) Guide to best practices for ocean acidification measurements. *PICES Special Publication* 3, 191 pp.
- His E, Seaman MNL, Beiras R (1997) A simplification of the bivalve embryogenesis and larval development bioassay method for water quality assessment. *Water Research* 31: 351–355.
- Bos OG, Philippart CJM, Cadée GC, van der Meer J (2006) Recruitment variation in *Macoma balthica* (L.): the match/mismatch hypothesis. *Marine Ecology Progress Series* 320: 207–214.
- Bos OG, Philippart CJM, van der Meer J (2007) Effects of temporary food limitation on development and mortality of *Macoma balthica* larvae. *Marine Ecology Progress Series* 330: 155–162.
- Conover WJ (1980) *Practical Nonparametric Statistics*, Second Edition, New York: John Wiley & Sons, Inc.
- Havenhand JN, Butler FR, Thorndyke MC, Williamson JE (2008) Near-future levels of ocean acidification reduce fertilization success in a sea urchin. *Current Biology* 18: R651–R652.
- Suwa R, Morita M, Iguchi A, Nakamura M, Shimada K, et al. (2010) Ocean acidification reduces sperm flagellar motility in broadcast spawning reef invertebrates. *Zygote* 18: 103–107.
- Gould MC, Stephano JL (2003) Polyspermy prevention in marine invertebrates. *Microscopy Research and Technique* 61: 379–388.
- Reuter KE, Lotterhos KE, Crim RN, Thompson CA, Harley CDG (2011) Elevated pCO₂ increases sperm limitation and risk of polyspermy in the red sea urchin *Strongylocentrotus franciscanus*. *Global Change Biology* 17: 163–171.
- Johnson CH, Epel D (1981) Intracellular Ph of Sea-Urchin Eggs Measured by the Dimethylxazolinedione (Dmo) Method. *Journal of Cell Biology* 89: 284–291.
- Parker LM, Ross PM, O'Connor WA (2010) Comparing the effect of elevated pCO₂ and temperature on the fertilization and early development of two species of oysters. *Marine Biology* 157: 2435–2452.
- Havenhand JN, Schlegel P (2009) Near-future levels of ocean acidification do not affect sperm motility and fertilization kinetics in the oyster *Crassostrea gigas*. *Biogeosciences* 6: 3009–3015.
- Van Damme S, Struyf E, Maris T, Ysebaert T, Dehairs F, et al. (1995) Spatial and temporal patterns of water quality along the estuarine salinity gradient of the Scheldt estuary (Belgium and the Netherlands): results of an integrated monitoring approach. *Hydrobiologia* 540: 26–45.
- Talmage SC, Gobler CJ (2009) The effects of elevated carbon dioxide concentrations on the metamorphosis, size, and survival of larval hard clams (*Mercenaria mercenaria*), bay scallops (*Argopecten irradians*), and Eastern oysters (*Crassostrea virginica*). *Limnology and Oceanography* 54: 2072–2080.
- Thorson G (1950) Reproductive and larval ecology of marine bottom invertebrates. *Biological Reviews* 25: 1–45.

53. Pineda J, Reynolds NB, Starczak VR (2009) Complexity and simplification in understanding recruitment in benthic populations. *Population ecology* 51: 17–32.
54. Reise K (1983) Biotic enrichment of intertidal sediments by experimental aggregates of the deposit-feeding bivalve *Macoma balthica*. *Marine Ecology Progress Series* 12: 229–236.